

ANTIMICROBIALS



TETRACYCLINES

BACTERIOSTATIC

ANTIMICROBIALS

TETRACYCLINES

Tetracycline

Oxytetracycline

Chlortetracycline

Doxycycline



ANTIMICROBIALS

Tetracyclines - Mechanism of Action

Inhibit aminoacyl-transfer-RNA
binding to 30S ribosomal subunit-
mRNA

ANTIMICROBIALS

Tetracyclines - Pharmacokinetics

Amphoteric- improves distribution

Lipid soluble - well distributed (+ milk)

Bioavailability - about 50% of oral dose

Liver metabolism & enterohepatic circulation

- tetracycline, oxytetracycline urine/bile
- Doxycycline - bile and faecal elimination

ANTIMICROBIALS

Tetracyclines

Chelated by divalent cations (Ca^{+2})

Penicillins and tetracyclines are antagonistic!!!
(except uterine boluses)

Intramuscular - irritating, painful Polyvinyl
pyrrolidone (PVP) causes less injury and
pain at IM site \$\$

ANTIMICROBIALS

Tetracyclines - toxicity

GIT- fatal enterocolitis/horse

- Diarrhoea, superinfections
- Pseudomembraneous colitis

Vitamin B deficiency

Yellow teeth and bones (antianabolic)

Contraindicated in pregnant animals

ANTIMICROBIALS

Tetracyclines - toxicity

Bitter - salivation

Cardiovascular collapse (IV)

Drug fever, rashes, photosensitivity

"Fanconi-like" syndrome (out of date)

Inhibit hepatic metabolism

Nephrotoxicity with large overdose

ANTIMICROBIALS

Tetracyclines - Spectrum of Activity

Broad spectrum antibiotics

Gram + and Gram -

Mycoplasma, Rickettsia

Chlamydia, protozoa

Tetracyclines

Gram Positive Bacteria

Staphylococcus, Strep, Clostridium

Listeria monocytogenes,

Anaerobes

Some efficacy but not first choice

Gram Negative Bacteria

Some efficacy – especially doxycycline e.g. Bordetella, Brucella, Pasteurella, Shigella

Actinobacillus ligniersi, Moraxella

Other susceptible:

Haemobartonella felis

Mycoplasma, Chlamydia

Leptospira - doxycycline


ANTIMICROBIALS TETRACYCLINES - USES

- Broad spectrum in activity but resistance limits use
- Pneumonia
- Uterine boluses
- footrot
- leptospirosis (doxycycline)
- black leg



ANTIMICROBIALS TETRACYCLINES - USES

- chlamydia (*Chlamydia felis* in cats)



ANTIMICROBIALS

Chloramphenicol & Florfenicol

Bacteriostatic

Activity (similar to tetracyclines):

Gram + / gram - (not *Pseudomonas*)

Rickettsia

Chlamydia (not DoC for *C felis* in cat)

ANTIMICROBIALS

Chloramphenicol - Pharmacokinetics

Highly lipid soluble (penetrates eye)

Bioavailability 100%

Excreted primarily in the urine

Accumulation in cat plasma

due to poor ability to glucuronidate

ANTIMICROBIALS

Chloramphenicol

Toxicity

Associated with aplastic anaemia

Banned in food producing animals

Warn owners if dispensing

ANTIMICROBIALS

Chloramphenicol

USES

Topical eye treatment for chlamydia

Good penetration to the eye & CNS

ANTIMICROBIALS

Florfenicol


Adverse effects - associated with
testicular atrophy in bulls

USES:

Respiratory diseases of cattle

Pink eye

Foot rot



ANTIMICROBIALS

Macrolides

Bacterostatic

Inhibit Protein Synthesis


ANTIMICROBIALS
MACROLIDES

- ◆ erythromycin
- ◆ azithromycin
- ◆ tylosin



ANTIMICROBIALS
MACROLIDES

- ◆ tilmicosin*
 - ◆ *toxic to humans
- ◆ spiramycin



ANTIMICROBIALS
MACROLIDES

- ◆ Bacteriostatic at usual doses
- ◆ Distributes well to most tissues
- ◆ Food interferes with oral use
- ◆ Liver metabolism
- ◆ Biliary excretion - primary
- ◆ Increased GIT motility (erythro)

ANTIMICROBIALS

MACROLIDES - Toxicity

- ◆ Muscle paralysis with anaesthetics
- ◆ FATAL DIARRHOEA!
 - ◆ Horses and small mammals
- ◆ Cardiac effects
 - ◆ Horses, pig and primates
- ◆ IM injections are painful

ANTIMICROBIALS

Macrolides and Lincosamides

Narrow spectrum Gram +

BUT ALSO:

Pasteurella

Bacteroides

Mycoplasma

Rickettsia

ANTIMICROBIALS

MACROLIDES - USES

- ◆ Small animal:
 - ◆ Campylobacter infections
 - ◆ Staphylococcus alternative
 - ◆ Mycoplasma pneumonia
- ◆ Tend to cause GI upset - erythromycin


ANTIMICROBIALS
MACROLIDES - USES

- ◆ Foals - Rhodococcus infections
- ◆ Large animal:
 - ◆ Pneumonia
 - ◆ Footrot
 - ◆ Mastitis
- ◆ Where bacteria are resistant to other antibiotics

ANTIMICROBIALS
LINCOSAMIDES

Bacterostatic

- ◆ Clindamycin
- ◆ Lincomycin



ANTIMICROBIALS
Lincosamides

Toxicity

Reactions at injection sites

GIT upset (do not use in horses or small mammals)

ANTIMICROBIALS
Lincosamides

Pharmacokinetics:

Basic drugs

Poor bioavailability (PO)

Good distribution

Enterohepatic circulation

ANTIMICROBIALS

LINCOSAMIDES

- ◆ Biotransformed by liver
- ◆ Half life is increased by liver and/or kidney disease

ANTIMICROBIALS

LINCOSAMIDES

- ◆ Excellent penetration into abscesses
- ◆ Upper respiratory tract infections
- ◆ Osteomyelitis
- ◆ Deep pyodermas

ANTIMICROBIALS

Macrolides and Lincosamides

USES:

Pyodermas

Pasteurellosis

Respiratory infections in pigs

Osteomyelitis

Clindamycin - toxoplasmosis

ANTIMICROBIALS

Tetracyclines, Chloramphenicol, Macrolides and Lincosamides

SUMMARY

Spectrum of Activity

Adverse reactions/toxicity

Uses

